

IUHET-378
December 1997

Precision W -Boson and Higgs Boson Mass Determinations at Muon Colliders¹

M. S. Berger

*Indiana University
Bloomington Indiana 47405*

Abstract. Precise determinations of the masses of the W boson and of the top quark could stringently test the radiative structure of the Standard Model (SM) or provide evidence for new physics. We analyze the excellent prospects at a muon collider for measuring M_W and m_t in the W^+W^- and ZH threshold regions. With an integrated luminosity of 10 (100) fb^{-1} , the W -boson mass could be measured to a precision of 20 (6) MeV, provided that theoretical and experimental systematics are understood. A measurement of $\Delta M_W = 6$ MeV would constrain the mass of a ~ 100 GeV Higgs to about ± 10 GeV. We demonstrate that a measurement at future colliders of the Bjorken process. With an integrated luminosity of 100 fb^{-1} it is possible to measure the Standard Model Higgs mass to within 45 MeV at a $\mu^+\mu^-$ collider for $m_h = 100$ GeV.

INTRODUCTION

Muon colliders offer a wide range of opportunities for exploring physics within and beyond the Standard Model (SM). An important potential application of these machines is the precision measurement of particle masses, widths and couplings. We estimate here the accuracy with which the W and Higgs boson masses can be determined from W^+W^- and ZH threshold measurements at a muon collider [1,2].

M_W MEASUREMENT AT THE $\mu^+\mu^- \rightarrow W^+W^-$ THRESHOLD

A muon collider is particularly well suited to the threshold measurement because the energy of the beam has a very narrow spread. The threshold cross section [3,4] is most sensitive to M_W just above $\sqrt{s} = 2M_W$, but a tradeoff exists between maximizing the signal rate and the sensitivity of the cross section to M_W . Detailed

¹⁾ Presented at the Workshop on Physics at the First Muon Collider and at the Front End of a Muon Collider, November 6-9, 1997, Fermi National Accelerator Laboratory.

analysis [5] shows that if the background level is small and systematic uncertainties in efficiencies are not important, then the optimal measurement of M_W is obtained by collecting data at a single energy

$$\sqrt{s} \sim 2M_W + 0.5 \text{ GeV} \sim 161 \text{ GeV},$$

where the threshold cross section is sharply rising.




FIGURE 1. The cross section for $\mu^+\mu^- \rightarrow W^+W^-$ in the threshold region for $M_W = 80.3$ GeV (solid) and $M_W = 80.1, 80.5$ GeV (dashed). The inlaid graph shows the region of the threshold curve where the statistical sensitivity to M_W is maximized. Effects of ISR have been included.

For a LEP2 measurement with 100 pb^{-1} of integrated luminosity the background and systematic uncertainties are, in fact, sufficiently small that the error for M_W will be limited by the statistical uncertainty of the measurement at $\sqrt{s} = 161$ GeV. But, at a muon collider at high luminosity, systematic errors arising from uncertainties in the background level and the detection/triggering efficiencies will be dominant unless some of the luminosity is devoted to measuring the level of the background (which automatically includes somewhat similar efficiencies) at an energy below the W^+W^- threshold. Then, assuming that efficiencies for the background and W^+W^- signal are sufficiently well understood that systematic uncertainties effectively cancel in the ratio of the above-threshold to the below-threshold rates, a very accurate M_W determination becomes possible.

The dominant background derives from $e^+e^- \rightarrow (Z/\gamma)(Z/\gamma)$ which is essentially energy independent [5] below 180 GeV. For our present analysis we model the background as energy independent, and accordingly assume that one measurement at an energy in the range 140 to 150 GeV suffices to determine the background.

We analyze our ability to determine the W mass via just two measurements: one at center of mass energy $\sqrt{s} = 161$ GeV, just above threshold, and one at $\sqrt{s} = 150$ GeV. The optimal M_W measurement is obtained by expending about

two-thirds of the luminosity at $\sqrt{s} = 161$ GeV and one-third at $\sqrt{s} = 150$ GeV. Combining the three modes, an overall precision of

$$\Delta M_W = 6 \text{ MeV} \quad (1)$$

should be achievable.

The combination of the measurements of the masses M_Z , M_W and m_t to such high precision has dramatic implications for the indirect prediction of the mass of the Higgs boson and for other sources of physics beyond the Standard Model. This is illustrated in Fig. 2. Assuming the current central values of M_W , m_t , $\alpha(M_Z)$ and $\alpha_s(M_Z)$, and that $L = 10 \text{ fb}^{-1}$ (100 fb^{-1}) is devoted to the measurement of m_t (M_W), the mass of the SM Higgs boson would be determined to be 260 GeV with an error of about ± 5 GeV from $\Delta m_t = 200$ MeV at a fixed M_W , and about ± 20 GeV from $\Delta M_W = 6$ MeV at a fixed m_t . For $m_h = 100$ GeV, the corresponding values would be ± 2 GeV and ± 10 GeV, respectively. More generally, the Δm_h value scales roughly like m_h .




FIGURE 2. Correlation between M_W and m_t in the SM with QCD and electroweak corrections for $m_h = 100, 300$ and 1000 GeV. The data point and error bars illustrate the possible accuracy for the indirect m_h determination assuming $M_W = 80.356 \pm 0.006$ GeV and $m_t = 175 \pm 0.2$ GeV. The widths of the bands indicate the uncertainty in $\alpha(M_Z)$.

HIGGS BOSON MEASUREMENT AT THE $\mu^+\mu^- \rightarrow Zh$ THRESHOLD

A very accurate determination of m_h is obtained by measuring the threshold cross section for the Bjorken Higgs-strahlung process [6] $\ell^+\ell^- \rightarrow Zh$. With integrated luminosity $L = 100 \text{ fb}^{-1}$, a 1σ precision of order 45 MeV is possible for a SM Higgs $m_h = 100$ GeV at a $\mu^+\mu^-$ collider. This error in m_h is smaller than that achievable via final state mass reconstruction for a typical detector, and would then be the most accurate determination of m_h at an e^+e^- collider.

The SM Higgs boson is easily discovered in the Zh production mode by running the machine well above threshold, *e.g.* at $\sqrt{s} = 500$ GeV. For $m_h \lesssim 2M_W$ the dominant Higgs boson decay is to $b\bar{b}$ and most backgrounds can be eliminated by b -tagging. The best means for measuring m_h will be to first determine m_h to within a few hundred MeV in $\sqrt{s} = 500$ GeV running, which will also yield a precise measurement of $\sigma(Zh)$, and then reconfigure the collider for maximal luminosity in the threshold energy region $\sqrt{s} \approx M_Z + m_h$.

In Fig. 3 we show the cross section for the Bjorken process $\ell^+\ell^- \rightarrow Zh$ for Higgs masses from 50 to 150 GeV. Since the threshold behavior is S -wave, the rise in the cross section (which is a few tenths of a pb) in the threshold region is rapid.




FIGURE 3. The cross section vs. $\sqrt{s}$ for the process $\mu^+\mu^- \rightarrow Z^*h \rightarrow f\bar{f}h$ for a range of Higgs masses.

In the ideal case that the normalization of the measured Zh cross section as a function of $\sqrt{s}$ can be precisely predicted, including efficiencies and systematic effects, sensitivity to the SM Higgs boson mass is maximized by a single measurement of the cross section at $\sqrt{s} = M_Z + m_h + 0.5$ GeV, just above the real particle threshold. As an example of the precision that might be achieved, suppose $m_h = 100$ GeV and backgrounds are neglected. The Zh cross section is 120 fb and is rising at a rate of 0.05 fb/MeV. With $L = 50 \text{ fb}^{-1}$ and including an overall (b -tagging, geometric and event identification) efficiency of 40%, this yields 2.4×10^3 events, or a measurement of the cross section to about 2%. From the slope of the cross section one concludes that a m_h measurement with accuracy of roughly 50 MeV is possible.

For a more precise estimate of the accuracy with which m_h can be measured, we employ b -tagging and cuts in order to reduce the background to a very low level. These cuts and other systematic uncertainties are discussed in more detail in Ref. [2]. The background is very much smaller than the signal unless m_h is close to M_Z . We note that electroweak radiative corrections to the cross section are estimated to be less than 1% for $m_H \sim 100$ GeV [10,11]. A precision of the SM

Higgs mass determination to within 45 MeV for $m_h = 100$ GeV may be achievable at a muon collider.

Outside the Standard Model the cross section generally depends on the Higgs mass, the ZZH coupling (g_{ZZh}) and the total Higgs width (Γ_H). In order to simultaneously determine m_h , g_{ZZh} and Γ_H , measurements could be made at the three c.m. energies $\sqrt{s} = m_h + M_Z + 20$ GeV, $\sqrt{s} = m_h + M_Z + 0.5$ GeV, and $\sqrt{s} = m_h + M_Z - 2$ GeV. The solid curves in Fig. 4 show the statistical precision that can be obtained for $m_h = 100$ GeV in a three-parameter fit to m_h , $g_{ZZh}^2 B(h \rightarrow b\bar{b})$ and Γ_H , including smearing effects from bremsstrahlung, beamstrahlung and beam energy spread at a muon collider, using an integrated luminosity of $100/3 \text{ fb}^{-1}$ at each of the above three values of $\sqrt{s}$ in the threshold region. The crosses in the center of the ellipses indicate the input values. With a three-parameter fit, the attainable error in m_h is about 110 MeV at the 1σ level. The second panel in Fig. 4 shows that there is significant sensitivity to the Higgs width Γ_H if it is of order 100 MeV. If Γ_H is very narrow (~ 3 MeV is predicted in the SM) then $\Delta m_h \sim \pm 80$ MeV is possible (see the dashed ellipse) from a fit to m_h and $\sigma(Zh)B(h \rightarrow b\bar{b})$ by devoting 50 fb^{-1} at each of two c.m. energies, $\sqrt{s} = m_H + M_Z + 20$ GeV and $\sqrt{s} = m_H + M_Z + 0.5$ GeV. Measurements that would simultaneously determine m_h , $\sigma(Zh)B(h \rightarrow b\bar{b})$ and Γ_H could be done at a level of accuracy that could distinguish a Standard Model Higgs boson from its many possible (*e.g.* supersymmetric) extensions.




FIGURE 4. Solid curves show the $\Delta\chi^2 = 1$ contours for determining the Higgs mass versus $g_{ZZh}^2 B(h \rightarrow b\bar{b})$, or versus Γ_H , by devoting $100/3 \text{ fb}^{-1}$ to each of the c.m. energies $\sqrt{s} = M_Z + m_h + 0.5$ GeV, $\sqrt{s} = M_Z + m_h + 20$ GeV and $\sqrt{s} = M_Z + m_h - 2$ GeV at a muon collider; b -tagging and cuts 1)–4) are imposed and initial state radiation and beam energy smearing are included. A Higgs mass $m_H = 100$ GeV is assumed. The dashed curve shows the $\Delta\chi^2 = 1$ contour that results when Γ_H is negligibly small and 50 fb^{-1} is devoted to each of the c.m. energies $\sqrt{s} = m_H + M_Z + 0.5$ GeV and $\sqrt{s} = m_H + M_Z + 20$ GeV.

CONCLUSION

A muon collider offers an unparalleled opportunity for precision W and Higgs mass measurements in the respective threshold regions. These measurements, however, require a collider that can deliver substantial luminosity.

ACKNOWLEDGMENTS

I thank V. Barger, J. F. Gunion and T. Han for a pleasant collaboration on the issues reported here. This work was supported in part by the U.S. Department of Energy under Grant No. DE-FG02-91ER40661.

REFERENCES

1. V. Barger, M.S. Berger, J.F. Gunion and T. Han, Phys. Rev. **D56**, 1714 (1997).
2. V. Barger, M. S. Berger, J. F. Gunion and T. Han, Phys. Rev. Lett. **78**, 3991 (1997).
3. T. Muta, R. Najima and S. Wakaizumi, Mod. Phys. Lett. **A1**, 203 (1986).
4. V. S. Fadin, V. A. Khoze and A. D. Martin, Phys. Lett. **B311**, 311 (1993); V. S. Fadin, V. A. Khoze, A. D. Martin and W. J. Stirling, Phys. Lett. **B363**, 112 (1995); V. S. Fadin, V. A. Khoze, A. D. Martin and A. Chapovsky, Phys. Rev. **D52**, 1377 (1995).
5. Z. Kunszt and W.J. Stirling *et al.*, hep-ph/9602352, in *Proceedings of the Workshop on Physics at LEP2*, eds. G. Alterelli, T. Sjostrand and F. Zwirner, CERN Yellow Report CERN-96-01 (1996), Vol. 1, p. 141; W.J. Stirling, Nucl. Phys. **B456**, 3 (1995).
6. J.D. Bjorken, *Proceedings of the Summer Institute on Particle Physics*, ed. M. Zipf (Stanford, 1976).
7. V. Barger, M.S. Berger, J.F. Gunion and T. Han, Phys. Rev. Lett. **75**, 1462 (1995); Phys. Rept. **286**, 1 (1997).
8. E. Boos, M. Sachwitz, H. J. Schreiber and S. Shichanin, Int. J. Mod. Phys. **A10**, 2067 (1995).
9. *Higgs Boson Discovery and Properties*, J.F. Gunion *et al.*, *Proceedings of the 1996 Snowmass Workshop*.
10. B.A. Kniehl, Z. Phys. **C55**, 605 (1992).
11. R. Hempfling and B. Kniehl, Z. Phys. **C59**, 263 (1993).